

Preservation of iron(II) by carbon-rich matrices in a hydrothermal plume

Brandy M. Toner^{1*}†, Sirine C. Fakra², Steven J. Manganini¹, Cara M. Santelli¹, Matthew A. Marcus², James W. Moffett³, Olivier Rouxel¹, Christopher R. German¹ and Katrina J. Edwards^{1,3}

Hydrothermal venting associated with mid-ocean ridge volcanism is globally widespread¹. This venting is responsible for a dissolved iron flux to the ocean that is approximately equal to that associated with continental riverine runoff². For hydrothermal fluxes, it has long been assumed that most of the iron entering the oceans is precipitated in inorganic forms. However, the possibility of globally significant fluxes of iron escaping these mass precipitation events and entering open-ocean cycles is now being debated³, and two recent studies suggest that dissolved organic ligands might influence the fate of hydrothermally vented metals^{4,5}. Here we present spectromicroscopic measurements of iron and carbon in hydrothermal plume particles at the East Pacific Rise mid-ocean ridge. We show that organic carbon-rich matrices, containing evenly dispersed iron(II)-rich materials, are pervasive in hydrothermal plume particles. The absence of discrete iron(II) particles suggests that the carbon and iron associate through sorption or complexation. We suggest that these carbon matrices stabilize iron(II) released from hydrothermal vents in the region, preventing its oxidation and/or precipitation as insoluble minerals. Our findings have implications for deep-sea biogeochemical cycling of iron, a widely recognized limiting nutrient in the oceans.

The mixing of hot, chemically reduced hydrothermal fluids with cold, oxygenated deep-sea water drives the dominant inorganic reactions of polymetallic sulphide precipitation and Fe oxidation and precipitation in plumes⁶. As a consequence of seawater entrainment into rising plumes, the entire volume of the global ocean must, on average, come into contact with hydrothermal fluids and particles in a relatively short time, 4000–8000 yr. Much of the trace element chemical reactivity observed in plumes is attributed, but has not been firmly linked, to the precipitation and surface reactivity of Fe oxyhydroxide minerals in both buoyant and neutrally buoyant plumes⁷. The rate of hydrothermal Fe(II) oxidation has been correlated with the redox state of local deep waters⁸; however, roles for dissolved organic carbon⁹ (DOC) and microbial activity¹⁰ have also been hypothesized.

Although hydrothermal plume processes influence important global ocean elemental cycles, there is little mechanistic information available about these processes owing to the dynamic character of the plume environment and the complexity of plume particles. Most knowledge of plume mineral composition and reactivity is inferred from either bulk digestion of filtered solids⁷ or scanning electron microscopy with elemental analysis⁶. Here we use synchrotron-based X-ray absorption spectromicroscopy to investigate Fe and C speciation at the

nanoscale in plume particles. We propose a mechanism for the production of an organic matrix capable of stabilizing hydrothermally vented Fe(II). Our results suggest that the assumption that Fe cycling in deep-sea hydrothermal plumes is driven solely by inorganic processes is no longer valid and that organic processes must also be considered.

Our spectromicroscopic observations and bulk chemical measurements indicate that organic C-rich matrices are pervasive in hydrothermal plume particles at the mid-ocean ridge (MOR) East Pacific Rise (EPR) 9° N at 2,504 m depth. The distribution and chemical speciation of Fe and C within small ($\leq 10 \mu\text{m}$ diameter) plume particles was examined by scanning transmission X-ray microscopy (STXM) and near-edge X-ray absorption fine-structure (NEXAFS) spectroscopy. The descending non-buoyant plume materials collected from Tica vent are very heterogeneous in size and composition, and contain high particulate organic carbon (POC) concentrations (for example, 6.7 wt%) (Figs 1a,e, 2a–c and Supplementary Fig. S1, Table 1). We observed C-bearing particles and aggregates consisting of a wide variety of biological debris (megafaunal detritus, Supplementary Fig. S1) and exopolymeric-like matrices with fibrous morphologies coating and aggregating particles (Figs 1f, 2a and Supplementary Fig. S6).

The presence of megafaunal detritus in plume-particle aggregates is consistent with the biologically active zone surrounding Tica vent in combination with the phenomenon of entrainment of near-vent sea water into the rising hydrothermal plume. However, the high organic C content and prevalence of exopolymer-like matrices in the vent particles was unexpected. In contrast to ultramafic-hosted hydrothermal venting, basalt-hosted vents such as those at EPR are not expected to produce large quantities of organic C through abiotic synthesis¹¹. In addition, deep-sea DOC is generally thought to be low in concentration and composed of fairly recalcitrant compounds. The exopolymer-like matrix observed here is composed of molecules that are quite labile—lipids, polysaccharides and proteins, as identified by C 1s NEXAFS spectroscopy (see Fig. 2b, Supplementary Fig. S6 and the Methods section)—and are presumably in equilibrium with a dissolved fraction of equally labile compounds. Carbon 1s NEXAFS spectra indicate that the POC is chemically heterogeneous at the nanometre scale, with composition consistent with mixtures of organic compounds (Fig. 2a,b and Supplementary Fig. S6). Consistent with a recent report on Fe(III)-complexing ligands in non-buoyant plumes at the Mid-Atlantic Ridge 5° S (ref. 4), the presence of a biologically labile and chemically complex pool of organic C in the vicinity of hydrothermal venting at the EPR has implications

¹Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA, ²Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA, ³Department of Biological Sciences, University of Southern California, Los Angeles, California 90089, USA. *Present address: Department of Soil, Water, and Climate, University of Minnesota—Twin Cities, St. Paul, Minnesota 55108, USA. †e-mail: toner@umn.edu.




Figure 1 | Iron spectromicroscopy of a Tica vent aggregate showing Fe speciation and its association with C. **a**, STXM image recorded at the Fe $2p_{3/2}$ edge (707.6 eV) showing three areas of interest (AOI) outlined in white. **b**, Iron NEXAFS spectra extracted from the three AOI presented with Fe(II) (pyrrhotite) and Fe(III) (ferrihydrite). Spectra from AOI 1 and 2 are multiplied by 4 and 2, respectively, for display purposes. **c–e**, Fe(III) (**c**), Fe(II) (**d**) and C (**e**) distribution maps. **f**, STXM image at the C 1s edge (300 eV). Scale bars are 1 μm .

for microbiological cycling and chemical speciation of important elements such as Fe.

The STXM images and Fe(II,III) maps shown in Fig. 1a,c–f demonstrate that Tica vent particles aggregate and these aggregates are mixtures of Fe(II) and Fe(III): they also indicate that Fe(II), and to a lesser extent Fe(III), is co-located with the background organic matrix. The small, optically dense particles distributed throughout this representative aggregate are Fe(III)-rich minerals (Fig. 1a, area 3). Although Fe $2p$ NEXAFS spectra cannot reveal the exact Fe(III) oxyhydroxide form, these minerals bear much spectral resemblance to Fe oxyhydroxides such as ferrihydrite (Fig. 1b) and goethite¹². Evenly dispersed Fe(II)-rich materials largely dominate Fe speciation in certain areas of the aggregate (Fig. 1a, area 2). The evenly dispersed Fe associated with the organic matrix is consistent with either: (1) Fe in aqueous solution, (2) Fe in sorption complexes or (3) Fe in a uniformly distributed precipitate with a particle diameter much less than 30 nm. We rule out dissolved Fe because the concentration would have to approach 0.1 mM to be detected by STXM–NEXAFS (ref. 13). We can also exclude uniformly distributed nanoparticulate minerals on the basis of Fe $2p$ NEXAFS spectra because the Fe(II) spectra (Fig. 1a, areas 1 and 2) do not match those of Fe(II)-sulphide, -carbonate and -silicate minerals (Fig. 1b and Supplementary Fig. S5). In other words, the Fe(II) measured in those areas is not present as extremely fine-grained pyrite, basalt glass or other typical Fe-bearing vent mineral fragments. Rather, the absence of discrete particles is consistent with a chemical association between Fe and C through sorption or complexation of Fe(II) to organic functional groups within the organic matrix. Therefore, the role of POC in Fe speciation could be analogous to aqueous Fe/organic ligand interactions.

Our characterization of the mineralogy of the Tica vent plume particles, using micro-focused X-ray diffraction and X-ray absorption spectroscopy, indicates the presence of Fe(III) oxyhydroxide minerals and a variety of sulphide minerals (see Supplementary Figs S2–S4). Although these observations are consistent with current thinking on the fate of hydrothermal Fe(II) in vent fluids, detection of Fe(II) in association with an organic matrix stands in stark contrast to this inorganic precipitation paradigm. To our knowledge, we are reporting the first observation of Fe(II) stabilized by POC in hydrothermal systems. Measurements of dissolved Fe(II) in the Arabian Sea oxygen minimum zone indicate that dissolved Fe(II) is only prevalent when dissolved oxygen concentrations are below 2 μM (ref. 14). However, our samples were collected from oxygenated ($\sim 100 \mu\text{M kg}^{-1}$) deep-ocean waters⁸ and stored in the presence of oxygen for ~ 6 months before analysis. The predicted half-life of Fe(II) at our sampling site is 3.5 h (ref. 8): suggesting that Fe(II) was stabilized against oxidation by molecular oxygen. We hypothesize that this stabilization may be due to complexation by organic matter produced by hydrothermal processes. Although many Fe-complexing organic compounds can enhance Fe oxidation rates, slow rates of Fe(II) oxidation in marine waters have been attributed to the presence of organic ligands¹⁵.

Strictly on the basis of available stability constant data for Fe(II)-complexing organic ligands, organo-sulphur compounds such as 2,3-dimercaptopropanol ($\text{C}_3\text{H}_8\text{OS}_2$), with two thiol (-SH) groups per molecule, are the best model candidates for Fe(II)-stabilizing functional groups: log K values are 15.8 and 28.0 for the mono and bis complexes, respectively¹⁶. This compound also complexes Fe(III) with stability constants approximately equal to those for Fe(II), and our data do support the presence of




Figure 2 | Carbon spectromicroscopy of a Tica vent aggregate showing biomolecule signatures and association with Fe. **a**, STXM image recorded at the C 1s edge (300 eV). **b**, Carbon NEXAFS spectrum extracted from area of interest 4 outlined in white (**c**) along with reference spectra from protein (bovine serum albumin), a Tica vent microbial cell (see Supplementary Fig. S1), polysaccharide (alginate), nucleic acid (DNA) and lipid (1,2-dipalmitoyl-*sn*-glycero-3-phosphoethanolamine). **c**, Carbon distribution map. **d**, Iron distribution map from the white square area of the carbon map. Scale bars are 1 μm for **a–c** and 500 nm for **d**.

Fe(III) associated with POC. We hypothesize that, at the EPR, organo-sulphur compounds form in the buoyant plume from mixing of H_2S -rich vent fluids with entrained organic-rich waters derived from the biologically active sea floor in the vicinity of the vent. Our hypothesis is also supported by the fact that chemical reactions between H_2S and DOC have been shown to produce aggregates of polysaccharides and organo-sulphur compounds in marine surface waters¹⁷. In addition, Sander *et al.*⁵ have already proposed the presence of metal-reactive thiols in deep-sea hydrothermal fluids. At the EPR 9° N, we may have the conditions necessary to produce flocculant organic materials and organo-S compounds: fluids have an excess of H_2S (S/Fe ratio of 2.38:1; ref. 18) and near-vent bottom waters contain high POC concentrations (Table 1). Our hypothesis is testable through dynamic laboratory experiments that mix EPR endmember-like fluids with EPR near-vent bottom sea

water or artificial sea water with specific DOC–POC characteristics. Furthermore, as nanoprobe extended X-ray absorption fine-structure spectroscopy beamlines are developed, Fe 1s measurements may provide extra information regarding the coordination environment of Fe(II) in the organic matrix.

Although few published data are available, several studies of DOC–POC in the vicinity of high-temperature venting provide evidence supporting our hypothesis that entrained, near-vent biological materials react with vent fluid constituents to form POC capable of stabilizing Fe(II) against oxidation. At the Endeavour ridge, greater than 95% of the C measured within the first 21 vertical meters of a buoyant hydrothermal plume was identified as having a near-bottom biological origin¹⁹. More recently, measurements of DOC at the Juan de Fuca ridge have demonstrated elevated concentrations in some biologically productive areas of diffuse venting²⁰. Dissolved organic ligands present in non-buoyant hydrothermal plumes at the Mid-Atlantic Ridge have been found to stabilize Fe(III) in solution (these complexes may represent 12–22% of the dissolved Fe in the deep ocean)⁴. Furthermore, organic ligands with thiol functionality and stability constants high enough to compete with the precipitation of sulphide minerals have been proposed for hydrothermal fluids⁵.

The possible sources of DOC–POC to hydrothermal plumes may include: (1) abiotic synthesis in endmember fluids; (2) entrainment of near-vent biological materials such as megafaunal detritus, larvae and microbial mat materials, as well as vertical fluxes from the overlying water²¹; and (3) the presence and activity of microorganisms through input of exudates and cellular materials. For basalt-hosted systems such as the EPR, the biologically productive near-vent environment is probably the dominant source of both POC, as observed here, and DOC (ref. 4) to hydrothermal plumes.

The organic matrix we observe in EPR descending plume particles has an exopolymeric-like appearance (Fig. 2a), and may fit mechanistically into the DOC–POC continuum proposed generally for marine systems^{22,23}. The EPR POC matrix bears strong spectral similarities to biological molecules—proteins, polysaccharides and lipids—which have also been identified, by wet chemical techniques, to form the building blocks of marine polymer gels elsewhere in coastal waters from Puget Sound and deep North Pacific and Arctic oceans²⁴. Furthermore, both lipids²⁵ and exopolymeric-like matrices with spherical morphologies²⁶ have been observed in hydrothermal plumes rising above the Juan de Fuca ridge, suggesting that our findings may be representative of general processes occurring at deep-sea hydrothermal vents.

The fate and transport of hydrothermal plume POC and its associated Fe is not known. However, long-distance transport of hydrothermally sourced particles from EPR has been demonstrated^{27–29}. If we assume that colloidal particles with similar chemical composition are present in the migrating plume, then the long-lived presence of Fe(II) observed in these particles provides

Table 1 | Summary of mass flux, selected elemental composition, POC* and PIC[†] for plume particles collected by sediment traps.

	Mass flux $\text{mg m}^{-2} \text{d}^{-1}$	Total Fe wt%	Total Al wt%	Total Mn wt%	POC* wt%	PIC [†] wt%
Average (21–23 June 2006) [‡]	10.8	2.69	0.67	0.08	6.70	4.99
Range (16 May–27 June 2006) [§]	8.3–29.9	2.69–10.6	0.67–1.29	0.08–0.45	6.39–10.6	0.84–4.99

*Particulate organic carbon.

[†]Particulate inorganic carbon.

[‡]Particles from this sediment trap collection bottle were used for mass flux and STXM measurements.

[§]Reported figures represent the full range of values measured during sediment trap deployment.

